

ON A CONNECTION BETWEEN NAIMARK'S DILATION THEOREM, SPECTRAL REPRESENTATIONS, AND CHARACTERISTIC FUNCTIONS

MISHKO MITKOVSKI

ABSTRACT. We give a Herglotz-type representation of an arbitrary generalized spectral measure. As an application, a new proof of the classical Naimark's dilation theorem is given. The same approach is used to describe the spectrum of all unitary rank-one perturbations of a given partial isometry.

1. INTRODUCTION

Let $\mathcal{B}$ be a family of all Borel sets on the unit circle $\mathbb{T}$. By a generalized spectral measure on $\mathbb{T}$ we mean a function $B : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{H})$ whose values are positive bounded self-adjoint operators on $\mathcal{H}$ such that $B(\emptyset) = 0$, $B(\mathbb{T}) = I$ and for every sequence $\Delta_1, \Delta_2, \dots$ of mutually disjoint Borel sets, we have

$$B(\Delta_1 \cup \Delta_2 \cup \dots) = \sum_{i=1}^{\infty} B(\Delta_i)$$

in the strong operator topology. If we require all the values to be orthogonal projections then we have an (ordinary) spectral measure. A classical theorem of Naimark [3] says that any generalized spectral measure can be represented as a projection of an ordinary spectral measure. This theorem is considered by many as the beginning of Dilation Theory. Since then many different proofs and generalization have appeared (e.g. [4, 6, 8, 7]). We propose yet another approach, which involves in a natural way characteristic functions and spectral representations of unitary operators; it also relates Naimark's theorem for the first time to the subject of rank-one perturbations of a given operator. The latter is another classical subject with a rich literature behind (see [5] and the references therein). The key idea in our approach is to obtain a representation for a generalized spectral measure which is reminiscent to the well known one:

$$\langle (U + zI)(U - zI)^{-1}h_1 | h_2 \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d \langle (E(\xi)h_1 | h_2 \rangle$$

relating a unitary operator U to its spectral measure E . We will show that if $\mathcal{K}$ is a closed subspace of $\mathcal{H}$ and the generalized spectral measure $B : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{K})$ is obtained from a spectral measure $E : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{H})$ by $B(\Delta) = P_{\mathcal{K}}E(\Delta)$ then the above mentioned representation is given by:

$$\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}k_1 | k_2 \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d \langle (B(\xi)k_1 | k_2 \rangle$$

where $\tilde{U} : U^*(\mathcal{K}) \rightarrow \mathcal{K}$ is the restriction of $P_{\mathcal{K}}U$ on $U^*(\mathcal{K})$ and $\Theta_S(z)$ is the characteristic function corresponding to the operator $S := (I - P_{\mathcal{K}})U$. In the case

when $\mathcal{K}$ is one-dimensional this relation reduces to a relation that D. Clark [2] used in his treatment of the rank-one perturbations of a restricted shift. This is the point where the connection between these seemingly unrelated concepts is made.

The paper is organized as follows. The next section presents some preliminary material. Section 3 contains the main theorem which contains the above mentioned representation. As an application, a new proof of Naimark's dilation theorem is given. In the last section an application to the rank-one perturbations of a partial isometry is given.

2. PRELIMINARIES

We recall the basic facts from the the model theory of completely nonunitary contractions developed by Sz. Nagy and Foias [6]. Let $T \in \mathcal{B}(\mathcal{H})$ be a contraction, $\|T\| \leq 1$. The self adjoint operators $D_T := (I - T^*T)^{1/2}$ and $D_{T^*} := (I - TT^*)^{1/2}$ are called the defect operators of T and the spaces $\mathcal{D}_T := \overline{D_T\mathcal{H}}$, $\mathcal{D}_{T^*} := \overline{D_{T^*}\mathcal{H}}$ are called the defect spaces for T . The defect indices are defined by $\partial_T := \dim \mathcal{D}_T$ and $\partial_{T^*} := \dim \mathcal{D}_{T^*}$. These indices measure, in a certain sense, how much a contraction differs from a unitary operator. If T is a partial isometry then the defect operators D_T and D_{T^*} are orthogonal projections onto the initial and the final space of T respectively.

The characteristic function of a contraction T is an operator-valued function $\Theta_T(z) : \mathcal{D}_T \rightarrow \mathcal{D}_{T^*}$ defined by:

$$\Theta_T(z) := -T + zD_{T^*}(I - zT^*)^{-1}D_T|_{\mathcal{D}_T}.$$

It is always an analytic contraction-valued function. A contraction-valued function $\Theta : \mathbb{D} \rightarrow \mathcal{B}(\mathcal{L}, \mathcal{L}^*)$ is called pure if $\|\Theta(0)l\| < \|l\|$ for $l \in \mathcal{L}$, $l \neq 0$. Two contraction-valued analytic functions $\Theta_1 : \mathbb{D} \rightarrow \mathcal{B}(\mathcal{L}_1, \mathcal{L}_1^*)$, $\Theta_2 : \mathbb{D} \rightarrow \mathcal{B}(\mathcal{L}_2, \mathcal{L}_2^*)$ are said to coincide if there exist unitaries $\omega : \mathcal{L}_1 \rightarrow \mathcal{L}_2$, $\omega_* : \mathcal{L}_1^* \rightarrow \mathcal{L}_2^*$ such that $\Theta_1(z) = \omega_*^{-1}\Theta_2(z)\omega$. There is an easy characterization of partial isometries in terms of their characteristic functions. Namely, a contraction T is a partial isometry if and only if $\Theta_T(0) = O$.

A contraction T is said to be completely nonunitary (c.n.u.) if it is not unitary on any of its invariant subspaces. There always exists a unique decomposition $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$ into subspaces reducing T , such that $T|_{\mathcal{H}_0}$ is unitary and $T|_{\mathcal{H}_1}$ is c.n.u..

One of the main theorems of Sz. Nagy and Foias theory is the fact that every c.n.u. contraction T on a separable Hilbert space is unitarily equivalent to an operator $\mathbf{T}$ acting on the space

$$\mathbf{H} = [H^2(\mathcal{D}_{T^*}) \oplus \overline{\Delta_T L^2(\mathcal{D}_T)}] \ominus \{\Theta_T u \oplus \Delta_T u : u \in H^2(\mathcal{D}_T)\},$$

by

$$\mathbf{T}(u \oplus v) = \mathbf{P}_{\mathbf{H}}(zu \oplus zv),$$

where $\mathbf{P}_{\mathbf{H}}$ is the orthogonal projection of $H^2(\mathcal{D}_{T^*}) \oplus \overline{\Delta_T L^2(\mathcal{D}_T)}$ onto $\mathbf{H}$ and $\Delta_T(t) := (I - \Theta_T^*(e^{it})\Theta_T(e^{it}))^{1/2}$. Here, as usual, $L^2(\mathcal{D}_T)$ is the space of $\mathcal{D}_T$ -valued square integrable functions on $\mathbb{T}$ and $H^2(\mathcal{D}_T)$ is the corresponding Hardy space.

The following fact will be important for us. For any given contraction-valued analytic function $\Theta : \mathbb{D} \rightarrow \mathcal{B}(\mathcal{L}, \mathcal{L}^*)$ which is pure, there exists a contraction $\mathbf{T}$ whose characteristic function coincides with $\Theta(z)$.

In the case when $\Theta_T(z)$ is an inner function (i.e. $\Theta_T(\zeta)$ is an isometry for a.e. $\zeta \in \mathbb{T}$) then $\mathbf{H}$ has a simpler form $\mathbf{H} = H^2(\mathcal{D}_{T^*}) \ominus \{\Theta_T u : u \in H^2(\mathcal{D}_T)\}$. This happens if and only if $T^{*n} \rightarrow O$ as $n \rightarrow \infty$. If, in addition, the defect indices of T are both equal to 1 then $\mathbf{H} = H^2(\mathbb{D}) \ominus \Theta_T H^2(\mathbb{D})$ is a space of analytic scalar-valued functions. In this case $\Theta_T(z)$ can be viewed as a usual (scalar-valued) inner function. In [2], all the unitary rank-one perturbations of this type of operators T are examined. It is shown there that all the rank-one unitary perturbations can be parametrized by points $\alpha \in \mathbb{T}$ and that the spectral measures σ_α for each of these operators can be easily obtained from the characteristic function $\Theta_T(z)$. Namely, they are determined by:

$$\frac{\alpha + \Theta_T(z)}{\alpha + \Theta_T(z)} = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\sigma_\alpha(\xi).$$

The collection of measures that are associated to a given inner function (or more generally to an analytic self-map of the disc) in this way are called Clark (or Aleksandrov-Clark) measures [5].

3. MAIN RESULTS

Let $U : \mathcal{H} \rightarrow \mathcal{H}$ be a unitary operator. Fix $h \in \mathcal{H}$. Then the function $\langle (U + zI)(U - zI)^{-1}h|h \rangle$ is holomorphic in $z \in \mathbb{D}$ with positive real part. Thus, there exists a unique measure μ_h on the unit circle $\mathbb{T}$ such that

$$\langle (U + zI)(U - zI)^{-1}h|h \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\mu_h(\xi).$$

Using polarization, for any $h_1, h_2 \in \mathcal{H}$ there exists a measure μ_{h_1, h_2} such that:

$$\langle (U + zI)(U - zI)^{-1}h_1|h_2 \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\mu_{h_1, h_2}(\xi).$$

Let Δ be any Borel set on the unit circle $\mathbb{T}$. Then $\mu_{h_1, h_2}(\Delta)$ is a skew-symmetric function of h_1 and h_2 , linear in h_1 , and bounded by $\|h_1\| \|h_2\|$. Therefore, it can be represented as

$$\mu_{h_1, h_2}(\Delta) = \langle E(\Delta)h_1|h_2 \rangle,$$

for some positive bounded operator $E(\Delta)$. It is well known that $E : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{H})$ is an ordinary spectral measure.

Let $\mathcal{K} \subset \mathcal{H}$ be a closed subspace and let $P_{\mathcal{K}} : \mathcal{H} \rightarrow \mathcal{K}$ be the orthogonal projection onto $\mathcal{K}$. Define $T, S : \mathcal{H} \rightarrow \mathcal{H}$ by $T = P_{\mathcal{K}}U$ and $S = (I - P_{\mathcal{K}})U$, respectively. Then T and S are partial isometries and the characteristic function $\Theta_S(z) : U^*(\mathcal{K}) \rightarrow \mathcal{K}$ satisfies $\Theta_S(0) = O$ (and hence is pure). It will be useful to denote by $\tilde{U}$ the restriction of U on the closed subspace $U^*(\mathcal{K})$, and view it as an operator from $U^*(\mathcal{K})$ to $\mathcal{K}$. Obviously, $\tilde{U}u = Tu = Uu$ for any vector $u \in U^*(\mathcal{K})$ and $\tilde{U}^*k = T^*k = U^*k$ for any $k \in \mathcal{K}$. Observe also that $\tilde{U}^*P_{\mathcal{K}} = T^*P_{\mathcal{K}} = U^*P_{\mathcal{K}}^2 = T^*$.

For fixed $k \in \mathcal{K}$, again $\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}k|k \rangle$ is a holomorphic function in $z \in \mathbb{D}$ with positive real part and consequently there exists a unique measure σ_k satisfying

$$\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}k|k \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\sigma_k(\xi).$$

By polarization again, for any $k_1, k_2 \in \mathcal{K}$ there exists a measure σ_{k_1, k_2} such that:

$$\left\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}k_1 | k_2 \right\rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\sigma_{k_1, k_2}(\xi).$$

For any Borel set Δ on the unit circle let $B(\Delta)$ be the positive self-adjoint operator satisfying

$$\sigma_{k_1, k_2}(\Delta) = \langle B(\Delta)k_1 | k_2 \rangle,$$

for all $k_1, k_2 \in \mathcal{K}$. It can be shown that $B : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{K})$ is a generalized spectral measure.

Theorem 3.1. *For any $k \in \mathcal{K}$ the following equality holds:*

$$\langle (U + zI)(U - zI)^{-1}k | k \rangle = \left\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}k | k \right\rangle.$$

Proof: The left- and right-hand side of the equality we are aiming to prove are equal to $2 \left\langle (I - \Theta_S(z)\tilde{U}^*)^{-1}k | k \right\rangle - \|k\|^2$ and $2 \left\langle (I - zU^*)^{-1}k | k \right\rangle - \|k\|^2$, respectively. Hence, it is equivalent to show that:

$$\left\langle (I - \Theta_S(z)\tilde{U}^*)^{-1}k | k \right\rangle = \left\langle (I - zU^*)^{-1}k | k \right\rangle.$$

To prove this equality, we first notice that

$$\begin{aligned} \left\langle (I - \Theta_S(z)\tilde{U}^*)^{-1}k | k \right\rangle &= \left\langle (I - zP_{\mathcal{K}}(I - zS^*)^{-1}\tilde{U}^*)^{-1}k | k \right\rangle \\ &= \left\langle (\tilde{U}\tilde{U}^* - zP_{\mathcal{K}}(I - zS^*)^{-1}\tilde{U}^*)^{-1}k | k \right\rangle \\ &= \left\langle \tilde{U}(\tilde{U} - zP_{\mathcal{K}}(I - zS^*)^{-1})^{-1}k | k \right\rangle \\ &= \left\langle \tilde{U}(\tilde{U}(I - zS^*)(I - zS^*)^{-1} - zP_{\mathcal{K}}(I - zS^*)^{-1})^{-1}k | k \right\rangle \\ &= \left\langle \tilde{U}(I - zS^*)(\tilde{U}(I - zS^*) - zP_{\mathcal{K}})^{-1}k | k \right\rangle \\ &= \left\langle \tilde{U}(I - zS^*)(\tilde{U}(I - zS^*) - z\tilde{U}^*P_{\mathcal{K}})^{-1}k | k \right\rangle \\ &= \left\langle \tilde{U}(I - zS^*)(I - zS^* - z\tilde{U}^*P_{\mathcal{K}})^{-1}\tilde{U}^*k | k \right\rangle. \end{aligned}$$

Now, recalling that $\tilde{U}^*P_{\mathcal{K}} = T^*$, $T^*k = \tilde{U}^*k$ we obtain

$$\begin{aligned} \left\langle (I - \Theta_S(z)\tilde{U}^*)^{-1}k | k \right\rangle &= \left\langle (I - zS^*)(I - zS^* - zT^*)^{-1}\tilde{U}^*k | \tilde{U}^*k \right\rangle \\ &= \left\langle (I - zS^* - zT^*)^{-1}T^*k | T^*k \right\rangle - \left\langle z(I - zS^* - zT^*)^{-1}T^*k | ST^*k \right\rangle \\ &= \left\langle (I - zS^* - zT^*)^{-1}T^*k | T^*k \right\rangle. \end{aligned}$$

Since $ST^*k = 0$, we finally have

$$\begin{aligned} \left\langle (I - \Theta_S(z)\tilde{U}^*)^{-1}k | k \right\rangle &= \left\langle (I - z(T + S)^*)^{-1}T^*k | T^*k \right\rangle \\ &= \left\langle (I - zU^*)^{-1}U^*k | U^*k \right\rangle \\ &= \left\langle U^*(I - zU^*)^{-1}k | U^*k \right\rangle \\ &= \left\langle (I - zU^*)^{-1}k | k \right\rangle. \end{aligned}$$

□

Remark 1. If $\mathcal{K} = \mathcal{H}$ then $S = O$ on $\mathcal{H}$ and hence $\Theta_S(z) = zI$. Also, clearly $\tilde{U} = U$. Thus, we can view $(\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}$ as a substitution for $(U + zI)(U - zI)^{-1}$ in the case when $\mathcal{K}$ is a true subspace.

Corollary 3.2. For any Borel set $\Delta \subset \mathbb{T}$, $B(\Delta) = P_{\mathcal{K}}E(\Delta)$.

Proof: To show the claim it is equivalent to show

$$\langle P_{\mathcal{K}}E(\Delta)k_1 | k_2 \rangle = \langle B(\Delta)k_1 | k_2 \rangle$$

for all $k_1, k_2 \in \mathcal{K}$. Since $P_{\mathcal{K}}$ is an orthogonal projection, this is equivalent to:

$$\langle E(\Delta)k_1 | k_2 \rangle = \langle B(\Delta)k_1 | k_2 \rangle.$$

Therefore, using polarization it suffices to prove

$$\langle E(\Delta)k | k \rangle = \langle B(\Delta)k | k \rangle$$

for all $k \in \mathcal{K}$. But, this easily follows from Theorem 3.1. □

Next we give a new proof of Naimark's dilation theorem.

Theorem 3.3 (Naimark [3]). Let $B : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{K})$ be a generalized spectral measure. Then there exist $\mathcal{H} \supset \mathcal{K}$ and an ordinary spectral family $E : \mathcal{B} \rightarrow \mathcal{B}(\mathcal{H})$ such that for any Borel set $\Delta \subset \mathbb{T}$, $B(\Delta) = P_{\mathcal{K}}E(\Delta)$.

Proof: Let $k_1, k_2 \in \mathcal{K}$. Define $\sigma_{k_1, k_2}(\Delta) = \langle B(\Delta)k_1 | k_2 \rangle$ for any Borel set $\Delta \subset \mathbb{T}$. Clearly, σ_{k_1, k_2} is a Borel measure on $\mathbb{T}$. For any $z \in \mathbb{D}$ define $F(z) : \mathcal{K} \rightarrow \mathcal{K}$ such that

$$\langle F(z)k_1 | k_2 \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\sigma_{k_1, k_2}(\xi)$$

for every $k_1, k_2 \in \mathcal{K}$. It is easy to see that $F(z)$ is a linear operator with a positive real part and $F(0) = I$. It is also analytic as an operator-valued function in $z \in \mathbb{D}$. The inverse $(I + F(z))^{-1}$ is defined on a dense subset of $\mathcal{K}$ since $F(z)$ has a positive real part. Set $\Theta(z) = (F(z) - I)(F(z) + I)^{-1}$ on that dense set. By continuity, $\Theta(z)$ can be extended on the whole $\mathcal{K}$ with $\|\Theta(z)\| \leq 1$. Moreover, $\Theta(z)$ is an analytic contraction-valued function with $\Theta(0) = O$ (and hence is pure). Thus, there exists a completely non-unitary contraction $S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ whose characteristic function $\Theta_S(z)$ coincides with $\Theta(z)$. More precisely, there exist unitary operators $\omega : \mathcal{D}_S \rightarrow \mathcal{K}$ and $\omega_* : \mathcal{D}_{S^*} \rightarrow \mathcal{K}$ such that $\Theta_S(z) = \omega_*^{-1}\Theta(z)\omega$. Define $\tilde{U} = \omega_*^{-1}\omega : \mathcal{D}_S \rightarrow \mathcal{D}_{S^*}$ and set $T : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ to be

$$Th = \begin{cases} \tilde{U}h, & h \in \mathcal{D}_S \\ 0, & h \in \text{Ker } \mathcal{D}_S. \end{cases}$$

Finally, define $U_1 = T + S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$. Clearly, U_1 is unitary. Let $E_1(\Delta)$ be the spectral measure corresponding to U_1 and let $B_1(\Delta)$ be the generalized spectral measure such that

$$\left\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}s_1 | s_2 \right\rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\langle B_1(\xi)s_1 | s_2 \rangle$$

for every $s_1, s_2 \in \mathcal{D}_{S^*}$. By the previous Corollary, for any Borel set Δ we have

$$B_1(\Delta) = P_{\mathcal{D}_{S^*}}E_1(\Delta).$$

Now, since

$$\left\langle (\tilde{U} + \Theta_S(z))(\tilde{U} - \Theta_S(z))^{-1}s_1 | s_2 \right\rangle = \left\langle (I + \Theta(z))(I - \Theta(z))^{-1}\omega_*s_1 | \omega_*s_2 \right\rangle$$

and $F(z) = (I + \Theta(z))(I - \Theta(z))^{-1}$, it follows that $B(\Delta) = \omega_* B_1(\Delta) \omega_*^{-1}$.

Define $\mathcal{H} = \mathcal{K} \oplus \text{Ker} D_{S^*}$ and $W_* = \omega_* \oplus I : \mathcal{H}_1 \rightarrow \mathcal{H}$. Clearly, W_* is unitary. Then $B(\Delta) = P_{\mathcal{K}} E(\Delta)$ where $P_{\mathcal{K}} = W_* P_{\mathcal{D}_{S^*}} W_*^{-1}$ and $E(\Delta) = W_* E_1(\Delta) W_*^{-1}$. One readily sees that $P_{\mathcal{K}} : \mathcal{H} \rightarrow \mathcal{H}$ defined this way is the orthogonal projection onto $\mathcal{K}$ and $E(\Delta)$ is a spectral measure on $\mathcal{H}$. $\square$

4. RANK-ONE PERTURBATIONS OF PARTIAL ISOMETRIES

Let $S : \mathcal{H} \rightarrow \mathcal{H}$ be a c.n.u. partial isometry with both defect indices equal to 1, i.e., $\partial_S = \partial_{S^*} = 1$. Denote by $\mathcal{K}$ and $\tilde{\mathcal{K}}$ the orthogonal complements of the final and the initial space of S , respectively. Fix two unit vectors $k \in \mathcal{K}$ and $\tilde{k} \in \tilde{\mathcal{K}}$. For any complex number α of modulus 1, define the linear operator U_α by

$$U_\alpha h = \begin{cases} Sh, & \text{if } h \perp \tilde{\mathcal{K}} \\ \alpha k, & \text{if } h = \tilde{k}. \end{cases}$$

Clearly, U_α are unitary operators; these are the only unitary rank-one perturbations of S . Consider $\mathcal{K}_\alpha := \overline{\text{span}}\{U_\alpha^n k : n \in \mathbb{Z}\}$. One can show by induction that for any $h \in \mathcal{K}_\alpha^\perp$ and $\beta \in \mathbb{T}$ we have that $U_\beta^n h = S^n h$ and $U_\beta^{*n} h = S^{*n} h$. Therefore, $\mathcal{K}_\beta^\perp \subset \mathcal{K}_\alpha^\perp$ and, by symmetry, $\mathcal{K}_\beta^\perp = \mathcal{K}_\alpha^\perp$. This space $\mathcal{K}_\alpha^\perp$ is reducing for S and S is unitary there. Since S is c.n.u., we have that $\mathcal{K}_\alpha = \mathcal{H}$ for all $\alpha \in \mathbb{T}$.

Next we will describe the spectrum of U_α . Let $\Theta_S(z)$ be the characteristic function of S . For each $z \in \mathbb{D}$, $\Theta_S(z)$ is an operator between two one-dimensional spaces $\tilde{\mathcal{K}}$ and $\mathcal{K}$. Denote by $\phi(z)$ the scalar valued analytic function such that $\Theta_S(z)\tilde{k} = \phi(z)k$ for each $z \in \mathbb{C}$. Clearly, $|\phi(z)| \leq 1$. Define also $\tilde{U}_\alpha : \tilde{\mathcal{K}} \rightarrow \mathcal{K}$ to be the operator sending $\tilde{k}$ to αk . Let σ_α be the measure on $\mathbb{T}$ for which

$$\langle (\tilde{U}_\alpha + \Theta_S(z))(\tilde{U}_\alpha - \Theta_S(z))^{-1} k | k \rangle = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\sigma_\alpha(\xi).$$

This is equivalent to

$$\frac{\alpha + \phi(z)}{\alpha - \phi(z)} = \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\sigma_\alpha(\xi),$$

which implies that σ_α is a Clark measure for $\phi(z)$. It is also immediate that the corresponding generalized spectral measure $B_\alpha(\Delta)$ is simply given by $B_\alpha(\Delta)k = \sigma_\alpha(\Delta)k$. If $E_\alpha(\Delta)$ is the spectral measure corresponding to U_α then it follows from Corollary 3.2 that

$$\langle E_\alpha(\Delta)k | k \rangle = \sigma_\alpha(\Delta).$$

Since $\overline{\text{span}}\{E_\alpha(\Delta)k : \Delta \text{ Borel subset of } \mathbb{T}\} = \mathcal{H}$ the last equality proves the following:

Corollary 4.1. *The spectrum of U_α coincides with the support of σ_α . Thus, it consists of the union of those points in $\mathbb{T}$ at which $\phi(z)$ cannot be analytically continued and those $\zeta \in \mathbb{T}$ at which $\phi(z)$ is analytically continuable with $\phi(\zeta) = \alpha$. The set of eigenvalues of U_α coincides with the set of all the atoms of σ_α .*

Remark 2. There are several well-known conditions describing the atoms of a Clark measure σ_α . An important one (goes back to M. Riesz) is the following: σ_α

has a point mass at ζ if and only if $\phi(z)$ has the nontangential limit α at ζ and for all (or one) β in $\mathbb{T}$ different from α ,

$$\int_{\mathbb{T}} \frac{d\sigma_{\beta}(\zeta)}{|\xi - \zeta|^2} < \infty.$$

Remark 3. In [2], a similar description (although with different methods) of the spectra is obtained for unitary rank-one perturbations of a restricted shift. Notice that not every partial isometry can be represented as a restricted shift on the space $H^2(\mathbb{D}) \ominus \Theta H^2(\mathbb{D})$ of scalar-valued functions. Thus, the proposition above is not implied by the results in [2].

Finally, we can also consider the more general case when S is c.n.u. with equal (possibly infinite) defect indices. Let again $\mathcal{K}$ and $\tilde{\mathcal{K}}$ be the orthogonal complements of the final and the initial space of S , respectively. For a unitary operator $A : \tilde{\mathcal{K}} \rightarrow \mathcal{K}$, similarly as in [1], we can define a unitary perturbation U_A of S by

$$U_A h = \begin{cases} Sh, & \text{if } h \perp \tilde{\mathcal{K}} \\ Ah, & \text{if } h \in \tilde{\mathcal{K}}. \end{cases}$$

As in the case of the rank-one perturbations, one can show that S c.n.u. implies $\overline{\text{span}}\{U_A^n k : k \in \mathcal{K}, n \in \mathbb{Z}\} = \mathcal{H}$. To describe the spectrum of U_A notice that by Theorem 3.1 we have

$$\begin{aligned} \langle (A + \Theta_S(z))(A - \Theta_S(z))^{-1} k | k \rangle &= \langle (\tilde{U}_A + \Theta_S(z))(\tilde{U}_A - \Theta_S(z))^{-1} k | k \rangle \\ &= \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d \langle E_A(\xi) k | k \rangle. \end{aligned}$$

Corollary 4.1 has the following analogue in this general case.

Corollary 4.2. *The spectrum of U_A consists of the union of those points in $\mathbb{T}$ at which $\Theta_S(z)$ cannot be analytically continued and those $\zeta \in \mathbb{T}$ at which $\Theta_S(z)$ is analytically continuable with $\Theta_S(\zeta) - A$ not invertible.*

Further extensions of these results will be considered in a subsequent paper.

REFERENCES

- [1] BALL, J. A. AND LUBIN, A. *On a class of contractive perturbations of restricted shifts*, Pacific J. Math., **63** (1976), 309-323.
- [2] CLARK, D. N. *One dimensional perturbations of restricted shifts*, J. Analyse Math., **25** (1972), 169-191.
- [3] NAIMARK, M. A. *Spectral functions of a symmetric operator*, Izv. Akad. Nauk. SSSR Ser. Mat., **4** (1940), 277-318.
- [4] NAIMARK, M. A. *On a representation of additive operator set functions*, Dokl. Acad. Sci. SSSR, **41** (1943), 373-375.
- [5] POLTORATSKI, A. AND SARASON, D. *Aleksandrov-Clark measures*, Recent advances in operator-related function theory, Contemp. Math., **393** (2006), 1-14.
- [6] SZ.-NAGY, B. AND FOIAS, C. *Harmonic analysis of operators on Hilbert space*, Translated from the French and revised, North-Holland Publishing Co., Amsterdam, 1970.
- [7] PAULSEN, V. *Completely Bounded Maps and Operator Algebras*, Cambridge University Press, 2003.
- [8] STINESPRING, W. F. *Positive Functions on C^* -algebras*, Proc. Amer. Math. Soc., **6** (1955), 211-216.

TEXAS A&M UNIVERSITY, DEPARTMENT OF MATHEMATICS, COLLEGE STATION, TX 77843,
USA
E-mail address: `mmitkov@math.tamu.edu`